

Classifying the Markets Volatility with ARMA Distance Measures

Edoardo Otranto

Dipartimento di Economia, Impresa e Regolamentazione,
Via Sardegna, 59, 07100 Sassari, Italy (*eotranto@uniss.it*)

Abstract

The financial time series are often characterized by similar volatility structures. The selection of series having a similar behavior could be important for the analysis of the transmission mechanisms of volatility and to forecast the time series, using the series with more similar structure. In this paper a metrics is developed in order to measure the distance between two GARCH models, extending well known results developed for the ARMA models. The statistic used to calculate it follows known distributions, so that it can be adopted as a test procedure. These tools can be used to develop an agglomerative algorithm in order to detect clusters of homogeneous series.

Keywords: GARCH models, clusters, agglomerative algorithm

1. Introduction

The financial time series are generally subject to co-movements and similar volatility structures, due to the strong influence among financial markets (see, for example, Bollerslev et al., 1994). Generally, “trouble” and “quiet” periods are transmitted from a market to another, but some markets absorb more these effects. The classification of financial time series in homogeneous clusters for similar volatility structures could be an important purpose for the financial analysts, also because movements in a given time series could help to forecast the movements of a similar time series.

In this paper we extend the distance measure proposed by Piccolo (1990) for AR models to the case of the GARCH (*Generalized AutoRegressive Conditional Heteroskedasticity*) family. As stressed by Otranto and Triacca (2002), this distance compares the stochastic properties of couples of series, or, in other words, the differences between the two data generating processes. In practice, the basic idea is that the estimation of GARCH models provides the statistical structure of the financial time series, so that the comparison of the models underlying the data generating processes is equivalent to compare the volatility structures of each series. The extension of this distance to the GARCH models is easy, considering the correspondence between GARCH and ARMA processes; in practice we express the residuals of a GARCH model in ARMA form and then we use, as in Otranto and Triacca (2002), the representation of ARMA models in AR terms (see, for example,

Brockwell and Davis, 1996) to apply the distance measure. This representation provides a formulation of the distance measure as a function of the GARCH parameters. In addition, the statistic calculated to measure the distance follows a known asymptotic distribution, so that it is possible to use it as a test procedure. If we select the series having distance not significantly different by zero, it is possible to cluster the homogeneous series. In particular, we develop an agglomerative algorithm, based on the distance measure proposed and on the results of the statistical test. The methodology is applied to classify the series of the returns of the main financial markets.

In the next section we will illustrate the instruments adopted to explicit the distance measure, with the study of the behavior of the distance proposed; we will pay a particular attention to the GARCH(1,1) model, which is the most popular model adopted for financial time series. Section 3 is devoted to the explanation of the use of this distance in classifying the volatility of markets; we develop an agglomerative algorithm and show an application of the procedure to nine stock exchange indices. Final remarks follow. In the final appendix, there is a report of some details on the AR metrics proposed by Piccolo (1990).

2. Distance between GARCH Models

The GARCH family is very popular in time series analysis and it is composed of a large set of models, which can represent different possible characteristics of financial time series; for a review of these models and their applications see Bollerslev et al. (1992) and Bollerslev et al. (1994).

For our purpose, we consider two time series following the models ($t = 1, \dots, T$):

$$\begin{aligned} y_{1,t} &= \mu_1 + \varepsilon_{1,t}, \\ y_{2,t} &= \mu_2 + \varepsilon_{2,t}; \end{aligned}$$

where $\varepsilon_{1,t}$ and $\varepsilon_{2,t}$ are mean zero heteroskedastic independent disturbances. In other terms, the two series have a constant mean, whereas the variances are time-varying. We suppose that the conditional variances $h_{1,t}$ and $h_{2,t}$ follow two different and independent GARCH(1,1) structures:

$$\begin{aligned} Var(y_{1,t}|I_{1,t-1}) &= h_{1,t} = \gamma_1 + \alpha_1 \varepsilon_{1,t-1}^2 + \beta_1 h_{1,t-1} \\ Var(y_{2,t}|I_{2,t-1}) &= h_{2,t} = \gamma_2 + \alpha_2 \varepsilon_{2,t-1}^2 + \beta_2 h_{2,t-1} \end{aligned} \quad (1)$$

where $I_{1,t}$ and $I_{2,t}$ represent the information available at time t and $\gamma_i > 0$, $0 < \alpha_i < 1$, $0 < \beta_i < 1$, $(\alpha_i + \beta_i) < 1$ ($i = 1, 2$). This is a typical representation for financial time series.

Equation (1) implies that the squared residuals follow ARMA(1,1) processes:

$$\varepsilon_{i,t}^2 = \gamma_i + (\alpha_i + \beta_i) \varepsilon_{i,t-1}^2 - \beta_i (\varepsilon_{i,t-1}^2 - h_{i,t-1}) + (\varepsilon_{i,t}^2 - h_{i,t}), \quad i = 1, 2 \quad (2)$$

where $\varepsilon_{i,t}^2 - h_{i,t}$ are mean zero errors, uncorrelated with past information. Substituting in

(2) the errors with their ARMA(1,1) expression, we obtain the AR(∞) representation:

$$\varepsilon_{i,t}^2 = \frac{\gamma_i}{1 - \beta_i} + \alpha_i \sum_{j=1}^{\infty} \beta_i^{j-1} \varepsilon_{i,t-j}^2 + (\varepsilon_{i,t}^2 - h_{i,t}). \quad (3)$$

In this form, the two GARCH(1,1) models can be compared in terms of the distance measure proposed by Piccolo (1990), explained in the final appendix. In particular, recalling that the general form of this metrics is:

$$\left[\sum_{j=1}^{\infty} (\pi_{1j} - \pi_{2j})^2 \right]^{1/2}, \quad (4)$$

where π_{1j} and π_{2j} are the autoregressive coefficients of two AR processes, using (3), we can express the distance between two GARCH(1,1) models as:

$$d = \left[\sum_{j=0}^{\infty} (\alpha_1 \beta_1^j - \alpha_2 \beta_2^j)^2 \right]^{1/2}.$$

Developing the expression in square brackets:

$$d = \left[\alpha_1^2 \sum_{j=0}^{\infty} \beta_1^{2j} + \alpha_2^2 \sum_{j=0}^{\infty} \beta_2^{2j} - 2\alpha_1 \alpha_2 \sum_{j=0}^{\infty} (\beta_1 \beta_2)^j \right]^{1/2} = \left[\frac{\alpha_1^2}{1 - \beta_1^2} + \frac{\alpha_2^2}{1 - \beta_2^2} - \frac{2\alpha_1 \alpha_2}{1 - \beta_1 \beta_2} \right]^{1/2} \quad (5)$$

Note that in the previous developments the constant $\gamma_i/(1 - \beta_i)$ was not considered; in effect, it does not affect the dynamics of the volatility of the two series, expressed by the autoregressive terms.

It is very simple to extend that to more general cases; in fact, the GARCH(p,q) model (Bollerslev, 1986):

$$h_t = \gamma + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_p \varepsilon_{t-p}^2 + \beta_1 h_{t-1} + \dots + \beta_q h_{t-q}$$

corresponds to the ARMA(p*,q) model, with $p^* = \max(p, q)$:

$$\varepsilon_t^2 = \gamma + (\alpha_1 + \beta_1) \varepsilon_{t-1}^2 + \dots + (\alpha_{p^*} + \beta_{p^*}) \varepsilon_{t-p^*}^2 - \beta_1 (\varepsilon_{t-1}^2 - h_{t-1}) - \dots - \beta_q (\varepsilon_{t-q}^2 - h_{t-q}) + (\varepsilon_t^2 - h_t).$$

Of course, if $p > q$, we put $\beta_{q+1} = \dots = \beta_p = 0$; if $q > p$, then $\alpha_{p+1} = \dots = \alpha_q = 0$.

The ARCH(p) model (Engle, 1982):

$$h_t = \gamma + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_p \varepsilon_{t-p}^2$$

corresponds to the AR(p) model:

$$\varepsilon_t^2 = \gamma + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_p \varepsilon_{t-p}^2 + (\varepsilon_t^2 - h_t);$$

the IGARCH(1,1) model (Engle and Bollerslev, 1986):

$$h_t = \gamma + (1 - \beta_1) \varepsilon_{t-1}^2 + \beta_1 h_{t-1}$$

corresponds to the IMA(1,1) model:

$$(\varepsilon_t^2 - \varepsilon_{t-1}^2) = \gamma - \beta_1 (\varepsilon_{t-1}^2 - h_{t-1}) + (\varepsilon_t^2 - h_t);$$

and so on.

In general, indicating with ϕ_k the generic AR coefficient and θ_j the generic MA coefficient of an ARMA model, we have:

$$\begin{aligned}\phi_k &= (\alpha_k + \beta_k), \\ \theta_j &= -\beta_j.\end{aligned}\tag{6}$$

To apply (4) we need the AR representation of the ARMA model; following Brockwell and Davis (1996), the iterative formula:

$$\pi_k + \sum_{j=1}^q \theta_j \pi_{k-j} = -\phi_k, \quad k = 0, 1, \dots$$

with $\phi_0 = 1$, can be applied. For the GARCH case, the previous relationship is equivalent to:

$$\pi_k = -(\alpha_k + \beta_k) + \sum_{j=1}^q \beta_j \pi_{k-j} = -\alpha_k + \sum_{j=1}^{q-1} \beta_j \pi_{k-j}.\tag{7}$$

Using (7) it is possible, applying (4), to compare every couple of GARCH models, not necessarily of the same order. In the remain of the work we will refer to GARCH(1,1) models, which are the most popular models for financial time series and for which the simple form (5) can be applied.

2.1 An Investigation about the GARCH(1,1) Distance

In this subsection we study more in detail the behavior of the distance (5), for various combination of the coefficients α_i and β_i . The behavior of the distance is clear when we pose $\beta_i = 0$ for $i = 1, 2$, which is the case of two ARCH(1) models. In this case, the distance shows a double linear dynamics, symmetric with respect to the point representing the equality of the two data generating processes. In Figure 1 the comparison of two ARCH(1) models, with coefficients varying in $[0.1, 0.9]$ with steps of 0.1 is shown; each line represents the distance between an ARCH(1) model with coefficient indicated in the box, and the ARCH(1) models with coefficients equal to the corresponding points on the horizontal axis.

When two GARCH(1,1) models are considered, the behavior is well different; in fact, for the contemporaneous presence of α_i and β_i , similar processes can seem different. In Figure 2 the behavior of the distance between two GARCH(1,1) models is shown; note that there is a vast region (approximately when $0.1 \leq \alpha_i \leq 0.3$ and $0.1 \leq \beta_i \leq 0.8$, for $i = 1, 2$) in which the distance is approximately zero. This would be clearer observing Figure 3, in which the dark spots indicate the zones in which the distance is not significantly different by zero in the case of $T = 3000$.¹ In addition, the equality of α_1 and α_2 or β_1 and β_2 cuts down the distance considerably; this is more evident observing the detail of three profiles in Figure 4.

¹ The test used depends from the coefficients of the GARCH models and the number of observations; it is described in the final appendix. We have used a large number of observations, generally being available large data sets for financial time series; increasing the number of observations, the spots will grow progressively narrower.

Figure 1: Distance between two ARCH(1) models




Figure 2: Distance between two GARCH(1,1) models




Figure 3: Regions with non significantly different by zero distance for GARCH(1,1) model and T=3000.





Figure 4: Profiles of three GARCH(1,1) distances



3. Clustering the Returns: An Agglomerative Algorithm

How could it be used in practical cases the distance developed in the previous section? The most obvious application is to create homogeneous groups having a similar volatility structure. For this purpose an usual agglomerative algorithm for cluster analyses could be used; it can be developed in the following steps:

1. choose an initial benchmark series;
2. insert in the group of the benchmark series all the series with a distance from it not significantly different by zero;
3. select the series with the minimum distance from the benchmark significantly different by zero; this series will be the new benchmark;
4. insert in the second group all the remaining series with a distance from the new benchmark not significantly different by zero;
5. repeat steps 3 and 4 until no series remain.

Note that, differently from the common cluster algorithms, in this case the number of groups is not fixed a priori or chosen after the clustering, but it derives automatically from the algorithm. Clearly, to classify the series we need a starting point, in the sense that the result will be different, changing the series adopted as initial benchmark. Alternatively, in applications with a small number of series, we can use each series as initial benchmark in different classifications and then verify if there are “strongest” structures.

In order to explain this algorithm, we consider the series of the returns of nine stock exchange indices from December 1, 1995 to February 5, 2001 (daily data, $T = 1352$); they refer to the following indices: CAC40 (*cac*), NIKKEI300 (*nik*), DAX30 (*dax*), SMI (*smi*), FTSE100 (*fts*), IBEX35I (*ibe*), DOW JONES (*dj*), BEL20 (*bel*), MIB30 (*mib*). First, a GARCH(1,1) model is estimated for each series, then the matrix of distances for each couple of series is calculated and finally the statistical test to verify the null of zero distance is applied. The estimations of coefficients are shown in Table 1, whereas the matrix of distances in Table 2.

In Table 3 are shown the results of the diagnostic test for each couple of indices (A indicates the case of acceptance of the null of distance 0, whereas R indicates the case of rejection).

In our example, using each series as initial benchmark, the 9 classifications provide three possible alternative distinct groups. Using as initial benchmark *cac*, *dax*, *fts*, *bel* and *mib*, the 2 groups obtained are formed by (*cac*, *nik*, *fts*) and (*dax*, *smi*, *ibe*, *dj*, *bel*, *mib*); using as initial benchmark *smi*, *ibe* and *dj*, the 2 groups are formed by (*cac*, *fts*) and (*nik*, *dax*, *smi*, *ibe*, *dj*, *bel*, *mib*); using *nik* as initial benchmark we separate (*dax*, *bel*, *mib*) from (*cac*, *nik*, *smi*, *fts*, *ibe*, *dj*). Combining the results we deduce that there are 2 strong groups, constituted by *cac* and *fts* on a hand and *dj*, *dax*, *smi* and *ibe* on the other hand. The *nik* stays in the middle, whereas *bel* and *mib* are very similar to the *dj* group, but distant from *nik*.

Table 1: GARCH(1,1) parameters estimation (standard errors in parentheses).

	cac	nik	dax	smi	fts
γ	0.0003 (0.0001)	0.0009 (0.0003)	0.0005 (0.0001)	0.0006 (0.0001)	0.0001 (3.86E-5)
α	0.0540 (0.0081)	0.0637 (0.0094)	0.0965 (0.0129)	0.0891 (0.0147)	0.0443 (0.0076)
β	0.9335 (0.0094)	0.9191 (0.0124)	0.8876 (0.0139)	0.8775 (0.0198)	0.9492 (0.0078)
	ibe	dj	bel	mib	
γ	0.0006 (0.0001)	0.0005 (0.0001)	0.0003 (8.91E-5)	0.0010 (0.0002)	
α	0.0838 (0.0109)	0.0829 (0.0089)	0.0989 (0.0130)	0.1159 (0.0179)	
β	0.8920 (0.0128)	0.8873 (0.0135)	0.8863 (0.0138)	0.8384 (0.0226)	

Table 2: Distances matrix.

	cac	nik	dax	smi	fts	ibe	dj	bel	mib
cac	0.000	0.019	0.076	0.063	0.022	0.054	0.053	0.080	0.101
nik	0.019	0.000	0.057	0.045	0.041	0.035	0.035	0.062	0.084
dax	0.076	0.057	0.000	0.025	0.097	0.024	0.030	0.004	0.041
smi	0.063	0.045	0.025	0.000	0.084	0.012	0.010	0.029	0.040
fts	0.022	0.041	0.097	0.084	0.000	0.076	0.075	0.101	0.121
ibe	0.054	0.035	0.024	0.012	0.076	0.000	0.007	0.029	0.050
dj	0.053	0.035	0.030	0.010	0.075	0.007	0.000	0.034	0.050
bel	0.080	0.062	0.004	0.029	0.101	0.029	0.034	0.000	0.040
mib	0.101	0.084	0.041	0.040	0.121	0.050	0.050	0.040	0.000

Table 3: Test results.

	cac	nik	dax	smi	fts	ibe	dj	bel	mib
cac		A	R	R	A	R	R	R	R
nik	A		R	A	A	A	A	R	R
dax	R	R		A	R	A	A	A	A
smi	R	A	A		R	A	A	A	A
fts	A	A	R	R		R	R	R	R
ibe	R	A	A	A	R		A	A	A
dj	R	A	A	A	R	A		A	A
bel	R	R	A	A	R	A	A		A
mib	R	R	A	A	R	A	A	A	

4. Concluding Remarks

In this paper an extension of the distance measure used to compare couples of ARMA models, developed by Piccolo (1990), is extended to the GARCH case. This extension avoids the possibility to group the financial series having a similar volatility structure and an agglomerative algorithm was developed to obtain homogeneous clusters. The final results of the algorithm depend on the series adopted as benchmark; anyway, this is not necessarily a weak point, because generally the behavior of the markets are evaluated with respect to a “dominant” market (for example, the U.S. stock exchange market, which influences the other markets or shares); on the other side, the detection of various clusters, obtained using as benchmark each market iteratively, will conduce probably to some “strong” form, or some interpretable behavior, as in the application of the previous section.

Clearly, the case of clustering is just a possible application of this instrument; another purpose could be to forecast assets, shares or stock exchange indices of the financial markets ; as well known, for the volatility transmission mechanisms, the information deriving from a market can influence the behavior of another market. Using the distance measure, it is possible to detect the most similar volatility structure for a certain series among a set of leading series, so that the knowledge of the latter could be used to forecast the volatility structure of the former.

Appendix: The AR Metrics

In this appendix there is a brief description of the AR metrics introduced by Piccolo (1990) and the considerations above its distribution developed in Corduas (1996) with extensions to the GARCH(1,1) case.

Let V_t be a zero-mean ARMA invertible process; then, it exists a sequence $\{\pi_j\}$ such that

$$\sum_{j=1}^{\infty} |\pi_j| < \infty$$

and

$$V_t = \sum_{j=1}^{\infty} \pi_j V_{t-j} + \varepsilon_t, \quad (8)$$

where ε_t is a white noise process with variance σ^2 .

Piccolo (1990) defines the distance between two ARMA invertible and independent processes V_{1t} and V_{2t} as

$$d = \left[\sum_{j=1}^{\infty} (\pi_{1j} - \pi_{2j})^2 \right]^{1/2}. \quad (9)$$

From (9) we have derived the GARCH(1,1) distance (5).

Piccolo (1989) shows that the asymptotic distribution of d^2 , given the independence hypothesis, is a linear combination of independent Chi-Square variables. In order to deal with the distance measure as a test procedure, Corduas (1996) proposes to approximate the distribution of d^2 with a single Chi-Square random variable. Under the null hypothesis:

$$\pi_{1j} = \pi_{2j} \quad \text{for each } j = 1, 2, \dots, \quad (10)$$

this distribution can be approximated with $a\chi_c^2 + b$, where χ_c^2 is a chi-squared random variable with c degree of freedom and, setting $t_i = \text{trace}(\tilde{\Sigma}^i)$:

$$a = t_3/t_2, \quad b = t_1 - t_2^2/t_3, \quad c = t_2^3/t_3^2. \quad (11)$$

This approximation has a good performance, as showed in Corduas (1996). In this case $\tilde{\Sigma} = \tilde{\Sigma}_1 + \tilde{\Sigma}_2$ and represents the covariance matrix of the AR coefficients in (9) under the null hypothesis. $\tilde{\Sigma}_1$ and $\tilde{\Sigma}_2$ represent respectively the estimated covariance matrices of the coefficients $\tilde{\pi}_1 = \{\pi_{1j}\}$ and $\tilde{\pi}_2 = \{\pi_{2j}\}$, obtained as functions of the maximum likelihood estimators of the parameters of the GARCH models, as showed in (7). For practical purposes, the vectors $\tilde{\pi}_1$ and $\tilde{\pi}_2$ will contain only the first k autoregressive coefficients of the representation (8), with k suitably high (in our applications we will use $k = 100$). The covariances matrices $\tilde{\Sigma}_1$ and $\tilde{\Sigma}_2$ can be obtained by:

$$\tilde{\Sigma}_i = \Gamma_i \tilde{\mathbf{V}}_i \Gamma_i',$$

where $\tilde{\mathbf{V}}_i$ is the covariance matrix of the estimated GARCH coefficients and Γ_i is a matrix containing the derivatives of the functions π_{ij} with respect the GARCH coefficients. For example, for the case of GARCH(1,1) model, the estimated parameters modeling the volatility structure will be $(\tilde{\alpha}_i, \tilde{\beta}_i)'$, whereas $\tilde{\pi}_i = (\tilde{\alpha}_i, \tilde{\alpha}_i \tilde{\beta}_i, \dots, \tilde{\alpha}_i \tilde{\beta}_i^{k-1})'$.

Note that, to map out Figure 3, we have not performed estimation procedures, having used the theoretical covariance matrix of ARMA(1,1) processes (Brockwell and Davis, 1996). For an ARMA(1,1) process with AR coefficient equal ϕ and MA coefficient equal θ , the covariance matrix is expressed by:

$$\mathbf{V}_{ARMA} = \frac{1 + \phi\theta}{T(\phi + \theta^2)} \begin{bmatrix} (1 - \phi^2)(1 + \phi\theta) & -(1 - \theta^2)(1 - \phi^2) \\ -(1 - \theta^2)(1 - \phi^2) & (1 - \theta^2)(1 + \phi\theta) \end{bmatrix} = c \begin{bmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{bmatrix};$$

taking into account (6), we obtain that:

$$\begin{aligned} Var(\alpha_i + \beta_i) &= Var(\alpha_i) + Var(\beta_i) + 2Cov(\alpha_i, \beta_i) = a_{11} \\ Var(\beta_i) &= a_{22} \\ Cov(\alpha_i + \beta_i, -\beta_i) &= -Cov(\alpha_i, \beta_i) - Var(\beta_i) = a_{12} \end{aligned}$$

As a consequence:

$$\mathbf{V}_{GARCH} = c \begin{bmatrix} (a_{11} + a_{22} + 2a_{12}) & -(a_{12} + a_{22}) \\ -(a_{12} + a_{22}) & a_{22} \end{bmatrix}.$$

In this way we can apply the test considering hypothetical GARCH(1,1) processes, without estimation step; the only sample information we need is the length of the series T .

Acknowledgments

I thank Umberto Triacca for our useful discussions.

References

- [1] Bollerslev T. (1986), Generalized Autoregressive Conditional Heteroskedasticity, *Journal of Econometrics*, 31, 307-321.
- [2] Bollerslev T., Chou R. Y., Kroner K. F. (1992): ARCH Modeling in Finance: A Review of the Theory and Empirical Evidence, *Journal of Econometrics*, 52, 5:59
- [3] Bollerslev T., Engle R. F., Nelson D. (1994), ARCH Models, in *Handbook of Econometrics, Vol. IV* (Engle R. F. and McFadden D. L. eds.), 2959-3038.
- [4] Brockwell, P.J., Davis, R. A. (1996): *Introduction to Time Series and Forecasting*, Springer-Verlag, New York: .
- [5] Corduas, M. (1996). Uno Studio sulla Distribuzione Asintotica della Metrica Autoregressiva, *Statistica*, 56, 321-332.
- [6] Engle R. F. (1982), Autoregressive Conditional Heteroskedasticity with Estimates of the Variance of U.K. Inflation, *Econometrica*, 50, 987-1008.
- [7] Engle R. F., Bollerslev T. (1986), Modelling the Persistence of Conditional Variances, in *Econometric Reviews*, 5, 1-50.
- [8] Otranto E., Triacca U. (2002), Measures to Evaluate the Discrepancy between Direct and Indirect Model-based Seasonal Adjustment, *Journal of Official Statistics*, 18, 511-530.
- [9] Piccolo D. (1989), On the Measure of Dissimilarity between ARIMA Models, *Proceedings of the A.S.A Meetings*, Business and Economic Statistics Sect., Washington D. C., 231-236.
- [10] Piccolo D. (1990), A Distance Measure for Classifying ARIMA Models, *Journal of Time Series Analysis*, 11, 153-164.