

On the stability of recursive least squares in the Gauss-markov model

Evens SALIES

University of Paris 1 - Panthéon Sorbonne.

e.salies@caramail.com

Problem and Motivation

Consider the classical model $\mathbf{y}_n = \mathbf{X}_n\boldsymbol{\beta} + \boldsymbol{\varepsilon}_n$ where \mathbf{X}_n is an $n \times p$ real matrix of fixed regressors, \mathbf{y}_n ($n \times 1$) a response vector, $\boldsymbol{\beta}$ is a $p \times 1$ vector of unknown coefficients, $\text{rk}(\mathbf{X}_n) = p$ for $n \geq p$. Let $\hat{\boldsymbol{\beta}}(n)$ denote the ordinary least squares estimate of $\boldsymbol{\beta}$ obtained from n observations, with $n \geq p$, and assume $\boldsymbol{\varepsilon}_n$ ($n \times 1$) is a vector of non-observable random disturbances with expectation $\mathbf{0}$ and variance $\sigma^2\mathbf{I}_n$.

An updating formula for $\hat{\boldsymbol{\beta}}(n+1)$ as a function of $\hat{\boldsymbol{\beta}}(n)$ is

$$\hat{\boldsymbol{\beta}}(n+1) - \boldsymbol{\beta} = \mathbf{W}^{-1}\mathbf{V}(\hat{\boldsymbol{\beta}}(n) - \boldsymbol{\beta}) + \mathbf{w}, \quad n = p, p+1, \dots \quad (1)$$

where $\mathbf{V} \equiv \mathbf{X}'_n\mathbf{X}_n$, $\mathbf{W} \equiv \mathbf{X}'_{n+1}\mathbf{X}_{n+1}$, $\mathbf{w} \equiv \mathbf{W}^{-1}\mathbf{x}\boldsymbol{\varepsilon}_{n+1}$, and \mathbf{x} denotes the vector of new observations at the values of the explanatory variables. Eq. (1) arises for example in Kalman filtering and recursive least squares theories, where the unknown $\boldsymbol{\beta}$ is considered as time-varying states of dynamic system (see the discussion in Kianifard and Swallow, 1996) and $\mathbf{W}^{-1}\mathbf{V}$ is often developed as $\mathbf{I}_p - (1+c)^{-1}\mathbf{V}^{-1}\mathbf{x}\mathbf{x}'$; c equals $\mathbf{x}'\mathbf{V}^{-1}\mathbf{x}$.

This exercise provides some properties of $\mathbf{W}^{-1}\mathbf{V}$, with all its eigenvalues and eigenvectors. Let $\mathbf{A} \equiv \mathbf{W}^{-1}\mathbf{V}$ have eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_p$. Show that

- (i) these eigenvalues are real, and that
- (ii) $\lambda_1 = 1/(1+c)$, $\lambda_2 = \lambda_3 = \dots = \lambda_p = 1$.

Solution and Discussion

(i) \mathbf{A} is the product between two real symmetric matrices. Let λ be an eigenvalue of \mathbf{A} , and $\mathbf{u} + i\mathbf{v}$ an associated eigenvector, where $i^2 = -1$. Then

$$\mathbf{A}(\mathbf{u} + i\mathbf{v}) = \lambda(\mathbf{u} + i\mathbf{v}).$$

Premultiplying both sides of this equation with \mathbf{W} leads to

$$\mathbf{V}(\mathbf{u} + i\mathbf{v}) = \lambda\mathbf{W}(\mathbf{u} + i\mathbf{v}).$$

As $\mathbf{W} = \mathbf{V} + \mathbf{x}\mathbf{x}'$ therefore the previous equation becomes

$$(1 - \lambda)\mathbf{V}(\mathbf{u} + i\mathbf{v}) = \lambda\mathbf{x}\mathbf{x}'(\mathbf{u} + i\mathbf{v}).$$

Premultiply both sides with $(\mathbf{u} - i\mathbf{v})'$. Because of the symmetry of \mathbf{V} we obtain

$$(1 - \lambda)(\mathbf{u}'\mathbf{V}\mathbf{u} + \mathbf{v}'\mathbf{V}\mathbf{v}) = \lambda((\mathbf{u}'\mathbf{x})^2 + (\mathbf{v}'\mathbf{x})^2).$$

This implies that λ is real.

□

(ii) The following determinant

$$\begin{aligned} |\mathbf{I}_p - \mathbf{A}| &= |\mathbf{I}_p - \mathbf{W}^{-1}\mathbf{V}| \\ &= |\mathbf{W}^{-1}(\mathbf{W} - \mathbf{V})| \\ &= |\mathbf{W}^{-1}| \cdot |\mathbf{x}\mathbf{x}'| \\ &= |\mathbf{W}|^{-1} \cdot 0 \\ &= 0 \end{aligned}$$

shows $\lambda = 1$ is a root of the characteristic equation $|\lambda\mathbf{I}_p - \mathbf{A}| = 0$. Now, let \mathbf{z} be an eigenvector of \mathbf{A} associated with the eigenvalue 1; therefore

$\mathbf{W}^{-1}\mathbf{V}\mathbf{z} = \mathbf{z}$ or $\mathbf{V}\mathbf{z} = \mathbf{W}\mathbf{z}$, which from the definition of \mathbf{W} implies

$$\underset{(p \times 1)}{\mathbf{0}} = \mathbf{x}\mathbf{x}'\mathbf{z},$$

showing \mathbf{z} is orthogonal to \mathbf{x} . Remaining eigenvalues of \mathbf{A} are given using Wolkowicz and Styan's inequalities. We need $\text{trace}(\mathbf{A})$.

$$\begin{aligned} \text{trace}(\mathbf{A}) &= \text{trace}(\mathbf{W}^{-1}\mathbf{V}) \\ &= \text{trace}(\mathbf{W}^{-1}(\mathbf{W} - \mathbf{x}\mathbf{x}')) \\ &= \text{trace}(\mathbf{I}_p - \mathbf{W}^{-1}\mathbf{x}\mathbf{x}') \\ &= p - \mathbf{x}'\mathbf{W}^{-1}\mathbf{x}. \end{aligned}$$

Moreover, premultiplying $\mathbf{W} = \mathbf{V} + \mathbf{x}\mathbf{x}'$ by $\mathbf{x}'\mathbf{W}^{-1}$ and postmultiplying it by $\mathbf{V}^{-1}\mathbf{x}$ implies $\mathbf{x}'\mathbf{W}^{-1}\mathbf{x} = c/(1+c)$. Consequently

$$\text{trace}(\mathbf{A}) = p - c/(1+c),$$

and it can be shown \mathbf{x} is an eigenvector of \mathbf{A} and $1/(1+c)$ the associated eigenvalue. Premultiplying \mathbf{A} with \mathbf{x}' gives

$$\begin{aligned} \mathbf{x}'\mathbf{A} &= \mathbf{x}'(\mathbf{I}_p - \mathbf{W}^{-1}\mathbf{x}\mathbf{x}') \\ &= \mathbf{x}' - (\mathbf{x}'\mathbf{W}^{-1}\mathbf{x})\mathbf{x}' \\ &= \left(1 - \frac{c}{1+c}\right)\mathbf{x}' \\ &= \frac{1}{1+c}\mathbf{x}'. \end{aligned}$$

As \mathbf{A} has real eigenvalues we can apply the inequalities of Wolkowicz and Styan reproduced in Magnus and Neudecker (1991, p. 239) to find the order of multiplicity of previously found eigenvalues:

$$\begin{aligned} m - s(p-1)^{1/2} &\leq \lambda_1 \leq m - \frac{s}{(p-1)^{1/2}} \\ m + \frac{s}{(p-1)^{1/2}} &\leq \lambda_p \leq m + s(p-1)^{1/2}, \end{aligned}$$

where $m = (1/p)\text{trace}(\mathbf{A})$ and $s^2 = (1/p)\text{trace}(\mathbf{A}^2) - m^2$.

We obtain

$$1/(1+c) \leq \lambda_1 \leq 1 - \frac{2}{p} \frac{c}{1+c} \quad (2)$$

$$1 \leq \lambda_p \leq 1 + \frac{(p-2)}{p} \frac{c}{1+c}. \quad (3)$$

From Theorem 4 in Magnus and Neudecker (1991, p. 203),

$$\begin{aligned} \lambda_1 &\leq \frac{\mathbf{x}'\mathbf{A}\mathbf{x}}{\mathbf{x}'\mathbf{x}} \leq \lambda_p \\ \Leftrightarrow \lambda_1 &\leq \frac{\mathbf{x}'(\mathbf{I}_p - \mathbf{W}^{-1}\mathbf{x}\mathbf{x}')\mathbf{x}}{\mathbf{x}'\mathbf{x}} \leq \lambda_p \\ \Leftrightarrow \lambda_1 &\leq 1 - \frac{\mathbf{x}'\mathbf{W}^{-1}\mathbf{x}}{\mathbf{x}'\mathbf{x}} \leq \lambda_p \\ \Leftrightarrow \lambda_1 &\leq 1 - \frac{c}{1+c} \leq \lambda_p \\ \Leftrightarrow \lambda_1 &\leq \frac{1}{1+c} \leq \lambda_p. \end{aligned}$$

Combination of Eq. (2) and this result gives $\lambda_1 = 1/(1+c)$, which implies equality holds on the left of Eq. (3), that is $\lambda_p = 1$ and the $p-1$ largest eigenvalues are equal (Magnus and Neudecker, 1991, p. 239).

□

References

HARVEY, A. C. (1990). *The Econometric analysis of time series*. 2nd ed., Cambridge, MA: MIT Press.

KIANIFARD F., and SWALLOW, H. (1996). A review of the development and application of recursive residuals in linear models. *Journal of the American Statistical Association*, 91(433), pp. 391-400.

MAGNUS, J. R. and NEUDECKER, H. (1991). *Matrix Differential Calculus with Applications in Statistics and Econometrics*. John Wiley. Printed in Great Britain.