

# A Note on Imposing Local Curvature in Generalized Leontief Models\*

Apostolos Serletis<sup>†</sup> and Asghar Shahmoradi  
Department of Economics  
University of Calgary  
Calgary, Alberta, T2N 1N4,  
Canada

September 12, 2005

---

\*Serletis gratefully acknowledges financial support from the Social Sciences and Humanities Research Council of Canada.

<sup>†</sup>Corresponding author. Phone: (403) 220-4092; Fax: (403) 282-5262; E-mail: Serletis@ucalgary.ca; Web: <http://econ.ucalgary.ca/serletis.htm>.

Running Head (40 characteres or less): ????????

Corresponding Author: Apostolos Serletis

Apostolos Serletis  
Department of Economics  
University of Calgary  
Calgary, Alberta, T2N 1N4  
Canada

Phone: (403) 220-4092  
Fax: (403) 282-5262  
E-mail: Serletis@ucalgary.ca  
Web: <http://econ.ucalgary.ca/serletis.htm>.

## Abstract

In this paper, we build on Ryan and Wales (1998) and Moschini (1999) and impose curvature conditions locally on the generalized Leontief model, introduced by Diewert (1974). In doing so, we exploit the Hessian matrix of second order derivatives of the reciprocal indirect utility function, unlike Ryan and Wales (1998) and Moschini (1999) who exploit the Slutsky matrix.

*JEL* classification: C3; C13; C51.

*Keywords*: Regularity; Slutsky matrix; Negative semidefinite.

# 1 Introduction

As Moschini (1999, p. 487) puts it “[a] critical implication of consumer-demand theory is that the expenditure function is concave in prices, which implies that the so-called Slutsky matrix is negative semidefinite. Yet this curvature property is often not maintained in empirical applications, and estimated demand functions frequently fail to satisfy the implications of a concave expenditure function.” For this reason, in the recent literature there is an attempt to treat the curvature property as a maintained hypothesis and built it into the models being estimated, very much like the homogeneity in prices and symmetry properties of neoclassical consumer theory.

In particular, Ryan and Wales (1998) suggest a relatively simple procedure for imposing local curvature conditions. Their procedure applies to those locally flexible demand systems for which, at the point of approximation, the  $n \times n$  Slutsky matrix  $\mathbf{S}$  can be written as

$$\mathbf{S} = \mathbf{B} + \mathbf{C} \tag{1}$$

where  $\mathbf{B}$  is an  $n \times n$  symmetric matrix, containing the same number of independent elements as the Slutsky matrix, and  $\mathbf{C}$  is an  $n \times n$  matrix whose elements are functions of the other parameters of the system. Curvature requires the Slutsky matrix to be negative semidefinite. Ryan and Wales (1998) draw on related work by Lau (1978) and Diewert and Wales (1987) and impose curvature by replacing  $\mathbf{S}$  in (1) with  $-\mathbf{K}\mathbf{K}'$ , where  $\mathbf{K}$  is an  $n \times n$  lower triangular matrix, so that  $-\mathbf{K}\mathbf{K}'$  is by construction a negative semidefinite matrix. Then solving explicitly for  $\mathbf{B}$  in terms of  $\mathbf{K}$  and  $\mathbf{C}$  yields

$$\mathbf{B} = -\mathbf{K}\mathbf{K}' - \mathbf{C},$$

meaning that the models can be reparameterized by estimating the parameters in  $\mathbf{K}$  and  $\mathbf{C}$  instead of the parameters in  $\mathbf{B}$  and  $\mathbf{C}$ . That is, we can replace the elements of  $\mathbf{B}$  in the estimating equations by the elements of  $\mathbf{K}$  and the other parameters of the model, thus ensuring that  $\mathbf{S}$  is negative semidefinite at the point of approximation, which could be any data point.

Ryan and Wales (1998) applied their procedure to three locally flexible functional forms — the almost ideal demand system of Deaton and Muellbauer (1980), the normalized quadratic of Diewert and Wales (1988), and the linear translog. Moreover, Moschini (1999) suggested a possible reparameterization of the basic translog of Christensen, Jorgenson, and Lau (1975) to

overcome some problems noted by Ryan and Wales (1998) and also imposed curvature conditions locally in the basic translog.

In this paper, we build on Ryan and Wales (1998) and Moschini (1999) and impose curvature conditions locally on the generalized Leontief model, introduced by Diewert (1974). In doing so, we exploit the Hessian matrix of second order derivatives of the reciprocal indirect utility function, unlike Ryan and Wales (1998) and Moschini (1999) who exploit the Slutsky matrix. The rest of the paper is organized as follows. Section 2 briefly discusses the generalized Leontief demand system while Section 3 discusses our procedure for imposing local curvature conditions. The final section provides a brief conclusion.

## 2 The Generalized Leontief Model

The generalized Leontief (GL) reciprocal indirect utility function, introduced by Diewert (1974), is

$$h(\mathbf{v}) = \alpha_0 + \sum_{i=1}^n \alpha_i v_i^{1/2} + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \beta_{ij} v_i^{1/2} v_j^{1/2} \quad (2)$$

where  $\mathbf{B} \equiv [\beta_{ij}]$  is an  $n \times n$  symmetric matrix of parameters and  $\alpha_0$  and  $\alpha_i$  are other parameters, for a total of  $(n^2 + 3n + 2)/2$  parameters.

Using Roy's identity the share equations of the GL demand system can be written as

$$s_i = \frac{\alpha_i v_i^{1/2} + \sum_{j=1}^n \beta_{ij} v_i^{1/2} v_j^{1/2}}{\sum_{j=1}^n \alpha_j v_j^{1/2} + \sum_{k=1}^n \sum_{m=1}^n \beta_{km} v_k^{1/2} v_m^{1/2}}, \quad i = 1, \dots, n \quad (3)$$

Because the share equations are homogenous of degree zero in the parameters, one can follow Barnett and Lee (1985) and impose the following normalization in estimation

$$2 \sum_{i=1}^n \alpha_i + \sum_{i=1}^n \sum_{j=1}^n \beta_{ij} = 1 \quad (4)$$

### 3 Local Curvature for GL Models

Curvature of the reciprocal indirect utility function requires that the Hessian matrix is negative semidefinite. We can therefore impose local curvature (that is, at the reference point) by evaluating the Hessian terms of (2) at  $v_i^* = 1 \forall i$ , as follows

$$H_{ij} = -\delta_{ij} \left( \alpha_i + \sum_{j=1, j \neq i}^n \beta_{ij} \right) + (1 - \delta_{ij})\beta_{ij}$$

where  $\delta_{ij} = 1$  if  $i = j$  and 0 otherwise. By replacing  $\mathbf{H}$  by  $-\mathbf{K}\mathbf{K}'$ , where  $\mathbf{K}$  is an  $n \times n$  lower triangular matrix and  $\mathbf{K}'$  its transpose, the above can be written as

$$-(KK')_{ij} = -\delta_{ij} \left( \alpha_i + \sum_{j=1, j \neq i}^n \beta_{ij} \right) + (1 - \delta_{ij})\beta_{ij} \quad (5)$$

Solving for the  $a_i$  and  $\beta_{ij}$  terms as a function of the  $(KK')_{ij}$  terms we can get the restrictions that ensure the negative semidefiniteness of the Hessian matrix (without destroying the flexibility properties of (2), since the number of free parameters remains the same). In particular, when  $i \neq j$  (5) yields

$$\beta_{ij} = -(KK')_{ij} \quad (6)$$

and for  $i = j$  it yields

$$(KK')_{ii} = a_i + \sum_{j=1, j \neq i}^n \beta_{ij}$$

Substituting  $\beta_{ij}$  from (6) in the above equation

$$(KK')_{ii} = a_i - \sum_{j=1, j \neq i}^n (KK')_{ij}$$

or

$$a_i = (KK')_{ii} + \sum_{j=1, j \neq i}^n (KK')_{ij}$$

which after some rearrangement yields

$$a_i = \sum_{j=1}^n (KK')_{ij} \quad (7)$$

As an example, for the case of three goods (6) and (7) imply the following six restrictions

$$\begin{aligned} \beta_{12} &= -k_{11}k_{21} \\ \beta_{13} &= -k_{11}k_{31} \\ \beta_{23} &= -(k_{21}k_{31} + k_{22}k_{32}) \\ a_1 &= k_{11}^2 + k_{11}k_{21} + k_{11}k_{31} \\ a_2 &= k_{21}^2 + k_{22}^2 + k_{11}k_{21} + k_{21}k_{31} + k_{22}k_{32} \\ a_3 &= k_{31}^2 + k_{32}^2 + k_{33}^2 + k_{11}k_{31} + k_{21}k_{31} + k_{22}k_{32} \end{aligned}$$

where the  $k_{ij}$  terms are the elements of the replacement matrix  $\mathbf{K}$ .

## 4 Conclusion

We have illustrated a simple procedure for imposing curvature conditions locally in generalized Leontief models, building on earlier work in this area by Ryan and Wales (1998) and Moschini (1999). The procedure is rooted in the work by Lau (1978) and Diewert and Wales (1987), but unlike the Ryan and Wales (1998) and Moschini (1999) procedures that exploits the Slutsky matrix, our procedure exploits the Hessian matrix of second order derivatives of the reciprocal indirect utility function.

## References

- [1] Barnett, W.A. and Y.W.Lee. “The Global Properties of the Minflex Laurent, Generalized Leontief, and Translog Flexible Functional Forms.” *Econometrica* 53 (1985), 1421-1437. Reprinted in W.A. Barnett and J. Binner (eds), *Functional Structure and Approximation in Econometrics*. Elsevier, Amsterdam (2004), Chapter 4, pp. 79-97.
- [2] Christensen, L.R., D.W. Jorgenson, and L.J. Lau. “Transcendental Logarithmic Utility Functions.” *American Economic Review* 65 (1975), 367-383.
- [3] Deaton, A. and J. N. Muellbauer. “An Almost Ideal Demand System.” *American Economic Review* 70 (1980), 312-326.
- [4] Diewert, W. Erwin. “Applications of Duality Theory.” In *Frontiers of Quantitative Economics* (Vol. 2), eds. M.D. Intriligator and D.A. Kendrick, Amsterdam: North-Holland (1974), pp. 106-171.
- [5] Diewert, W. Erwin and Terence J. Wales. “Normalized Quadratic Systems of Consumer Demand Functions.” *Journal of Business and Economic Statistics* 6 (1988), 303-312.
- [6] Diewert, W. Erwin and Terence J. Wales. “Flexible Functional Forms and Global Curvature Conditions.” *Econometrica* 55 (1987), 43-68.
- [7] Lau, L.J. “Testing and Imposing Monotonicity, Convexity, and Quasi-Convexity Constraints.” In *Production Economics: A Dual Approach to Theory and Applications* (Vol. 1) M. Fuss and D. McFadden (eds.), Amsterdam: North Holland, (1978), p. 409-453.
- [8] Moschini, Giancarlo. “Imposing Local Curvature in Flexible Demand Systems.” *Journal of Business and Economic Statistics* 17 (1999), 487-490.
- [9] Ryan, David L. and Terence J. Wales. “A Simple Method for Imposing Local Curvature in Some Flexible Consumer-Demand Systems.” *Journal of Business and Economic Statistics* 16 (1998), 331-338.